

Description of Bulgarian System for Operational Pollution Forecast, version 3

Dimiter Syrakov¹, Maria Prodanova¹, Kiril Slavov¹, Igljka Etropolska¹,
Kostadin Ganev², Nikolai Miloshev², Todor Ljubenov³

¹*National Institute of Meteorology and Hydrology, Bulgarian Academy of Sciences, 66 Tzarigradsko Chausee, Sofia 1784, Bulgaria*

²*National Institute of Geophysics, Geodesy and Geography, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., Bl.3, Sofia 1113, Bulgaria*

³*Space Research and Technology Institute, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., Bl.1, Sofia 1113, Bulgaria*

INTRODUCTION

The Air Quality (AQ) is a key element for the well-being and quality of life of European citizens. According to the World Health Organization, air pollution severely affects the health of European citizens. There is increasing evidence for adverse effects of air pollution on the respiratory and the cardiovascular system as a result of both acute and chronic exposure. In particular, a significant reduction of life expectancy by a year or more is assumed to be linked to long-term exposure to high air concentrations of particulate matter (PM). There is considerable concern about impaired and detrimental air quality conditions over many areas in Europe, especially in urbanized areas, in spite of about 30 years of legislation and emission reduction. Current legislation, e.g. the Ozone daughter directive 2002/3/EC (European Parliament, 2002), requires informing the public on AQ, assessing air pollutant concentrations throughout the whole territory of Member States and indicating exceedances of limit and target values, forecasting potential exceedances and assessing possible emergency measures to abate exceedances. For the purpose, modeling tools must be used in parallel with air pollution measurements. The goals of reliable air quality forecasts are the efficient control and protection of population exposure as well as possible emission abatement measures. In last years the concept of “chemical weather” arises and in many countries respective forecast systems are being developed along with the usual meteorological weather forecasts (see, for instance, Sofiev et al., 2006, Poupkou et al., 2008, Monteiro et al., 2005, San Jose et al., 2006, and others).

Air pollution easily crosses national borders. It would be cost-effective and beneficial for citizens, society and decision-makers that national chemical weather forecast and information systems were networked across Europe. For the purpose several projects in the European Framework Programs (GEMS, PROMOTE, MEGAPOLI, MACC, PASODOBLE etc.) as well as the COST Action ES0602 “Towards a European Network on Chemical Weather Forecasting and Information Systems” were launched aiming at providing a forum for harmonizing, standardizing and benchmarking approaches and practices in data exchange and multi-model capabilities for air quality forecast and (near) real-time information systems in Europe. It is supposed to examine existing, and work out new solutions for integrating the development efforts at national and international levels. One can find several CW systems’ (performance and descriptions) in the Action’s web-portal (<http://www.chemicalweather.eu/Domains>).

Bulgaria joined the COST Action ES0602 from its very beginning. This participation invoked a project supported by the National Science Fund with the Bulgarian Ministry of Education, Youth and Science. Its main purpose was to create a Prototype of Bulgarian Chemical Weather Forecast and Information system (BgCWFIS, v.1) intended to provide timely, informative and reliable forecasts tailored to the needs of various users (see Syrakov et al. 2009, 2012). Later on, partly in the frame of EU FP7 project PASODOBLE (“Promote Air Quality Services Integrating Observations – Development of Basic Localized Information for Europe”) new versions of the system (v.2 and v.3) were elaborated with downscaling of the service to resolution of 1 km over

Sofia city. Here, the version 3 of BgCWFIS will be described and its end-user products will be presented.

MODELS USED

BgCWFIS is designed in a way to fit the real-time constraints and to deliver forecasts for the next days on an hourly basis. US EPA Models-3 air quality modeling system is used, consisting of:

- **CMAQ v.4.6** - Community Multi-scale Air Quality model, <http://www.cmaq-model.org/>, Denis et al. (1996), Byun and Ching (1999), Byun and Schere (2006), the Chemical Transport Model (CTM);
- **WRF v.3.2.1** - Weather Research and Forecasting Model, <http://www.wrf-model.org/>, Skamarock et al. (2005), the meteorological pre-processor to CMAQ;
- **SMOKE v.2.4** - Sparse Matrix Operator Kernel Emissions Modelling System, <http://www.smoke-model.org/>, Coats and Houyoux (1996), Houyoux and Vukovich (1999), CEP (2003), the emission pre-processor to CMAQ.

In the System, WRF is driven by the NCEP GFS (Global Forecast System) data that can be accessed freely from <http://www.ftp.ncep.noaa.gov/data/nccf/com/gfs/prod/>. This data is global weather forecast in GRIB-2 format with space resolution of $1^{\circ} \times 1^{\circ}$ and 6-hour time resolution. The downloading of this data is invoked automatically every day at 00:00Z. 84-hour runs starting at 12:00Z of the previous day are used; the first 12 hours of the period being spinning-up followed by a 3-day weather forecast. The chemical weather forecast duration is from 00:00Z of the current day to 00:00Z of the forth day after (3-day forecast).




Fig 1a. Downscaling the meteorological (WRF) forecast from domain Europe (81 km) to domain Bulgaria (09 km)




Fig 1b. Downscaling meteorological and chemical forecasts from domain Bulgaria (09 km) to domain Sofia city (01 km); in blue - WRF domains, in red - CMAQ domains.

MODEL DOMAINS

The nesting capabilities of WRF are used to downscale the meteorological forecasts from European region to Sofia-city area. The resolution of the mother domain (Europe) is 81 km, big enough in order to correspond to the GFS met-data resolution. Four other domains are nested in it and in each other – Balkan Peninsula (27 km resolution), Bulgaria (9 km), Sofia district (3 km) and

Sofia city (1 km) as shown in Fig. 1a,b. The decrease of space step from domain to domain is by factor of 3 – value recommended for both models.

In version 3 of BgCWFIS, WRF performs meteorological weather forecast for the all 5 domains while CMAQ performs chemical weather forecast over the last 3 domains (Bulgaria – Sofia region – Sofia city, Fig.1b). The chemical Boundary Conditions (BC) for Bulgaria domain are taken from the PASODOBLE joint airshed. Particularly, in this version of the System, data from SILAM model of Finish Meteorological Institute (FMI) is exploited (Sofiev et al., 2006, 2008, Vira and Sofiev, 2012, Kukkonen et al., 2012). SILAM forecast is 4 days long (to be extended to 5 days) that fits perfectly the operational schedule of BgCWFIS. Each day, after the meteorological processing, its performance continues with downloading 3-day data from SILAM yesterday's run that ensures full 3-day set of today's boundary conditions. The downloading exploits the web-based THREDDS system of SILAM (http://silam.fmi.fi/thredds/ncss/grid/sialm_european/runs/Slout, Slout = "silam_european_RUN_date&time") that permits to download only the necessary part of SILAM output. From the whole European domain of SILAM, forecast only for a small geographical region embracing Bulgaria (see Fig.2) is prepared by THREDDS as a single NetCDF file. This region is situated from 40° to 45° by latitude and from 21° to 30° by longitude. In addition to the region description, the request to THREDDS contains also the required pollutants, forecast periods etc. Special Linux script forming and executing the respective wget-command has been kindly provided by FMI. The invocation of this script delivers the necessary SILAM data in few minutes. This data is further processed by an interface program performing pollutants' mapping, horizontal and vertical interpolation etc. as to produce Bulgaria domain BC-file to CMAQ. The two minor domains take its BC from the senior ones.




Fig 2. Piece of SILAM forecast domain and CMAQ (domain Bulgaria) boundary points (red).

INFORMATION FLOW

The models indicated above are linked with a number of Linux scripts and interface programs in a way to be able to pre-calculate the future level of many air pollutants for every System's domain. The data flow diagram is presented in Fig.3.

The Models-3 elements are denoted by white boxes on the diagram, the blue boxes present the different kinds of input information. Those are: the meteorological numerical forecast that drives the meteorological pre-processor of the System (WRF) downloaded from NCEP's GFS data web-site; the CMAQ's Bulgaria domain BC downloaded from SILAM THREDDS system; the emission inventory data for 2005 provided by TNO, Netherlands (Denier van der Gon et al., 2010), gridded according to the 3 CMAQ's domains, and the respective three sets of land-use data (USGS data base, <http://landcover.usgs.gov/>). The land use data is extracted by WRF pre-processing module. The brown boxes present FORTRAN programs aimed at emission input modeling of Area Sources

(AS) and Large Point Sources (LPS), as well as data post-processing (extracting, archiving, image processing).




Fig 3. BgCWFIS, ver.3, information flow diagram. The white boxes present Models-3 elements, the blue ones – input data, and the brown ones – specific pre-and post-processing routines. The blue arrows denote the transfer of meteorological boundary conditions between the domains and the red arrows – the chemical boundary conditions. The black arrows show the exchange of information between the System's elements in each separate domain.

The data exchange in the System is denoted with arrows. The color arrows show the data exchange between computational domains, while the black arrows – the data exchange inside each one of the domains. Shortly, the WRF outputs feed MCIP (Meteorology-Chemistry Interface Processor) module of Models-3 system; MCIP prepares the meteorology input to CMAQ which is used also by SMOKE for calculating Biogenic Sources (BgS) emissions on the base of the respective LandUse data. The gridded inventory data (AInven, PInven) feeds AEmis and PEmis programs that produce the respective AS- and LPS-emission files. SMOKE is used once more to merge AS-, LPS- and BgS-data in a common emission input to CMAQ. Finally, the CMAQ output is post-processed in a way to extract the most important pollutants, to archive them, to produce hourly images with concentration distribution of 5 key pollutants (CO, NO₂, SO₂, Ozone and PM₁₀) as well as an Air Quality Index (AQI). In addition, a new interesting compliment to AQI is visualized – the dominant pollutant in the index. All images are uploaded to the respective web-site. These procedures are applied in the next order: WRF for all 5 domains of the system, for the last 3 domains - MCIP, AEmis, PEmis, SMOKE (BgS and merge), CMAQ and post-processing, consecutively. Meanwhile, WRF and CMAQ produce boundary condition files to be used by the inferior domains.

METEOROLOGICAL MODELLING

The Weather Research and Forecasting (WRF) Model is a next generation meso-scale numerical weather prediction system designed to serve both operational forecasting and atmospheric research needs. It is an evolutionary successor to the MM5 model. The creation and further development of WRF is due to the collaborative efforts of several US institutions like NCAR, NOAA, NCEP and

others. The WRF is a fully compressible and non-hydrostatic model with terrain-following hydrostatic pressure coordinate. The grid staggering is the Arakawa-C type. One can find more info on <http://www.wrf-model.org/index.php>. Free download can be made on http://www.mmm.ucar.edu/wrf/users/download/get_sources.htm.

In BgCWFIS, WRF-ARW (Advanced Research WRF), version 3.2.1, is exploited. The vertical structure is of 27 levels. The Analysis Nudging option (Four-Dimensional Data Assimilation, Stauffer and Seaman, 1994) is switched on for the first computational domain, only, i.e. WRF forecast is nudged to the meteorological driving data (GFS).

WRF-ARW offers multiple physics options that can be combined in any way. Here, well-tried schemes are used, namely:

Physics Options	Parameterization
Microphysics	WSM6 scheme (Hong and Lim, 2006)
Cumulus Parameterization	Kain-Fritsch scheme (Kain, 2004)
Planetary Boundary Layer	YSU scheme (Hong et al., 2006)
Longwave Radiation	RRTM scheme (Mlawer et al., 1997)
Shortwave Radiation	Dudhia scheme (Dudhia, 1989)
Land Surface Model	NOAH LSM scheme (Chen and Dudhia, 2001)

The coupling of a meteorological and chemical transport models is not a trivial issue. Since almost all meteorological models are not built for air quality modeling purposes, interface processing is needed. Such an element in Models-3 system is MCIP (Meteorology-Chemistry Interface Processor). Here, MCIP v.3.6. is applied. MCIP deals with issues related to data format translation, diagnostic estimations of parameters not provided by WRF (like dry deposition velocities for various species), extraction of data for appropriate window domains, and reconstruction of meteorological data on different grid and layer structures. As far as WRF-ARW and CMAQ are concerned, both utilize the Arakawa-C grid and conformal map projections and no special horizontal interpolation is required. MCIP may modify the vertical structure by interpolation between WRF layers. The sigma-levels of CMAQ in this application are 14:

$$\sigma = 1.0, 0.993, 0.99, 0.98, 0.965, 0.945, 0.91, 0.86, 0.8, 0.74, 0.65, 0.55, 0.4, 0.2, 0.0$$

EMISSION MODELLING

CMAQ demands its emission input in specific format reflecting the time evolution of all pollutants accounted for by the chemical mechanism used (CB-IV in this case). Emission inventories are used as row data for anthropogenic emission processing. The inventories are made on annual basis for big territories; many pollutants are estimated as groups (VOC and PM_{2.5} for instance). Preparation of emission input to a Chemical Transport Model requires emission processing. Such emission processing component in EPA Models-3 system is SMOKE but it is partly used, here, because its quite strong relation to US emission sources specifics. In BgCWFIS, SMOKE is used only for calculating BgS emissions and for merging AS-, LPS- and BgS-files into a CMAQ emission input file. The area source emissions and the large point source emissions are prepared by the interface programs AEmis and PEmis.

Inputs to AEmis and to PEmis are the respective gridded AS- and LPS-inventories. For the moment, TNO inventory for 2005 (Denier van der Gon et al., 2010) is exploited partly for Bulgaria domain, TNO being the Netherlands's Organization for Applied Scientific Research. For Bulgaria itself and for the other Bulgarian domains, the National inventory for 2010 as provided by Bulgarian Executive Environmental Agency is used. That means TNO inventory is used only for the territories outside Bulgaria in the mother CMAQ's domain.

The TNO produced several sets of inventories for different years. The anthropogenic sources in this inventories are distributed over 10 SNAPs (Selected Nomenclature for Air Pollution) classifying them according to the processes leading to harmful material release into the atmosphere (EMEP/CORINAIR, 2002). The 2005 TNO inventory has resolution of 0.125°×0.0625° (about 7×8

km). It is distributed as a comma- or tab-delimited text-file. Each line of the file contains data for a single box, namely the center of mesh coordinates, the country, the type of source (A/P), the SNAP, and the yearly emissions of 8 pollutants. The SNAP 7 (road transport) is presented as 5 sub-SNAPs. The pollutants are: methane (CH₄), carbon oxide (CO), nitric oxides (NO_x), sulfur oxides (SO_x), non-methane volatile organic compounds (NMVOC), ammonia (NH₃), Particulate Matter with $d < 10 \mu\text{m}$ (PM₁₀) and Particulate Matter with $d < 2.5 \mu\text{m}$ (PM_{2.5}).

As already stated, the inventory data can not be used directly as emission input to CTMs. Three operations must be applied to this data, preliminary: gridding, temporal allocation (imposing of time variations on the yearly values) and speciation (splitting of group pollutants to several simpler or lump pollutants).

The **gridding** is recalculation of the inventory data to the grids used. A web-based GIS system is created for the purpose. The TNO data is introduced in the system. Number of so-called custom grids can be defined on the base of the standard grid description (projection parameters, mean meridian, central point coordinates, distances to the grid origin, number of points, resolution). The system recalculates the inventory quantities for each cell, type of source and country.

Additional functionalities of the system are developed linked to the Bulgarian inventory. A specific feature of this inventory is that the data for SNAP 1-6 is attributed to particular sources with their coordinates. For SNAP 7-10 total country amounts of the released pollutants are available. First is the possibility of the GIS-system for aggregation of SNAPs 1-6 to a custom grid. In contrast, SNAP 7-10 inventory data must be disaggregated to the same grid using different geometric elements (surrogates). For instance, SNAP 7 (Road transport) is disaggregated by *HighWays* (15%), *RoadsClass1* (25%), *RoadsClass2* (20%) and *TownStreets* (40%). SNAP 8 (Non-road transport) uses *Airports* (45%), *RoadsClass2* (20%), *RailRoads* (25%), *AgricultureAreas* (10%). For SNAP 9 (Waste disposal) the surrogates are *Population* (50%) and *WasteAreas* (50%). Finally, SNAP 10 (Agriculture) is distributed according to *AgricultureAreas* (30%) and *Pastures* (70%). It is worth to mention that new surrogates can be added to the system and used in the disaggregation. The share of each surrogate can be changed, too.

The **temporal allocation** is made on the base of temporal profiles, provided by TNO (Bultjes et al., 2003). They are used also in both AQMEII exercises (<http://aqmeii.jrc.ec.europa.eu>). According to the anthropogenic activity the profiles are divided into three groups – Monthly, Weekly, and Hourly profiles. The first two are country-, SNAP- and pollutant-specific; the hourly factors refer to the local time and are SNAP-specific, only. Together with them, TNO provides a vertical profile for large point sources that is SNAP-specific.

The **speciation** profiles in this application are elaborated on the base of US EPA ones (<http://www.epa.gov/ttn/chief/emch/speciation/>) using the expert approach. Coincidence between main US sources with European SNAPs is obtained, first. The weighted averages of the respective speciation profiles are accepted as SNAP-specific splitting factors, weights being the percentage of contribution of every source type in total emission in particular SNAP. In such a way VOC, PM_{2.5}, NO_x and SO_x speciation profiles are derived. It must be noticed that the choice of US source types and the weights of their contribution to the respective SNAP emissions are quite subjective. Much more work is needed to elaborate country-specific profiles.

The gridded inventories are input to the emission processing programs AEmis and PEEmis, where the remaining two procedures are applied. In PEEmis, vertical allocation takes place as well. Both programs produce respective emission file. The AEmis output is 2-dimensional, the PEEmis one – 3-dimensional. Both outputs are in IO/API NetCDF format and contain hourly data for all forecast period.

The biogenic emissions are prepared by SMOKE by the BEIS-3.13 mechanism (Schwede et al., 2005) on the base of the gridded LandUse data. SMOKE merges the 3 emission files in a common CMAQ emission input – hourly data in NetCDF format.

OPERATIONAL PERFORMANCE OF BGCWFIS

Fourteen σ -levels with varying thickness determine the vertical structure of CMAQ. The Planetary Boundary Layer (PBL) is presented by the lowest 8 of these levels.

The CMAQ v.4.6 output consists of various files containing concentration, deposition, visibility and other variables. The concentration output is a NetCDF file with 3-D hourly data for 78 pollutants, of which: 52 gaseous, 21 aerosols (Aitken and accumulation modes), 5 aerosol distributions (3 by number, 2 by aerosol area).

The last box in Fig. 1 (upper-right) tags the post-processing that is quite important making BgCWFIS results visual. First of all the post-processing program XtrCON extracts part of the pollutants for archiving and further handling. Only surface values of the most important 17 pollutants are saved – 8 gases and 9 aerosols. Part of these pollutants is more or less monitored and they are referred in the European legislation with the respective thresholds. It must be mentioned that the sum of all aerosol compounds forms PM10 that is usually measured. The coarse particle matter (CPRM) is one of the saved aerosol compounds, so $PM2.5=PM10-CPRM$ is also calculated and archived as separate pollutant. All this data is stored in a respective archive file.

As to make the results of BgCWIS, ver.3, operation public, specialized web-site was created on the server <http://www.niggg.bas.bg/cw3/>. For the moment it presents 5 key pollutants (CO, NO₂, SO₂, Ozone, and PM10). In addition maps of the time-space distribution of an Air Quality Index (AQI) can be displayed.

In the moment, several AQI are exploited in various countries (Leeuw and Wim, 2005). Here, the UK index (p.25 of the just cited report) is chosen as one with best link to the calculated and visualized pollutant concentrations. The UK AQI presents air pollution level in 10 indexes and four bands according to the impact on the health of people who are sensitive to air pollution. In band “Low” (indexes 1-3) effects are unlikely to be noticed; in band “Moderate” (indexes 4-6) mild effects may be noticed amongst sensitive individuals, unlikely to require action; in band “High” (indexes 7-9) significant effects may be noticed by sensitive individuals and action to avoid or reduce these effects may be needed; in the last band “Very high” (index 10) the effects on sensitive individuals described for 'High' levels of pollution may worsen.

Each of the concentrations of the above 5 pollutants are classified in one of those 10 indexes, with boundaries given in Table 1. The overall index describes the total air pollution on the basis of the highest band for any individual pollutant.

Table 1. Boundaries between Index points for each pollutant

	Index	O ₃		NO ₂		SO ₂		CO		PM10	
		$\mu\text{g}/\text{m}^3$	ppb	$\mu\text{g}/\text{m}^3$	ppb	$\mu\text{g}/\text{m}^3$	ppb	mg/m ³	ppm	$\mu\text{g}/\text{m}^3$	TEOM
Low	1	0-32	0-16	0-95	0-49	0-88	0-32	0-3.8	0.0-3.2	0-21	0-16
	2	33-66	17-32	96-190	50-99	89-176	33-66	3.9-7.6	3.3-6.6	22-42	17-32
	3	67-99	33-49	191-286	100-149	177-265	67-99	7.7-11.5	6.7-9.9	43-64	33-49
Moderate	4	100-126	50-62	287-381	150-199	266-354	100-132	11.6-13.4	10.0-11.5	65-74	50-57
	5	127-152	63-76	382-477	200-249	355-442	133-166	13.5-15.4	11.6-13.2	75-86	58-66
	6	153-179	77-89	478-572	250-299	443-531	167-199	15.5-17.3	13.3-14.9	87-96	67-74
High	7	180-239	90-119	573-635	300-332	532-708	200-266	17.4-19.2	15.0-16.5	97-107	75-82
	8	240-299	120-149	636-700	333-366	709-886	267-332	19.3-21.2	16.6-18.2	108-118	83-91
	9	300-359	150-179	701-763	367-399	887-1063	333-399	21.3-23.1	18.3-19.9	119-129	92-99
Very High	10	360 or more	180 or more	764 or more	400 or more	1064 or more	400 or more	23.2 or more	20 or more	130 or more	100 or more

The described in Leeuw and Wim (2005, p.25) AQI, the time scale of concentrations are as follow: Ozone and NO₂ – hourly means, SO₂ – 15 minute means, CO – 8-hour running means, PM10 – 24-hour running means. In this application hourly means for all pollutants are used.

In addition to AQI, the dominant pollutant in this index (dpAQI) is determined and presented for each cell of the domains that possibly will be useful information for decision makers.

The software package PAVE (http://www.ie.unc.edu/cempd/EDSS/pave_doc/index.shtml) is used for visualization of the above 7 variables. PAVE supports its own meta-language that allows drawing plots in an automatic way by invoking the respective scripts.

In Fig.4, an example of BgCWFIS, ver.3, web-site is displayed. A particular pollutant or AQI is invoked by clicking in the list at the left side of the page. Note, that together with hourly ozone concentrations, two types of ozone daily maxima can be visualized.




Fig 4. Example of BgCWFIS, version 3, web-site.

The region of forecast can be chosen by clicking on one of the small images on the right. In the center of the page hourly concentration field of the chosen pollutant or AQI is situated. Putting the mouse cursor on one of the points from the point column on the right side of the image invokes the forecast field for the respective hour. Cursor placed over “Play” leads to animation of the forecast. Under each pollutant’s image, respective thresholds according to Bulgarian legislation (harmonized with European one) are shown. Under AQI images description of this index is presented. Here is the link to dpAQI images. At the bottom of pollutants list a link to a pdf-file with description of Bulgarian Chemical Weather Forecast and Information System, version 3, is placed.

CONCLUSION

The Bulgarian Chemical Weather Forecast and Information System, version 3, is designed on the base of US EPA Models-3 system: MM5 (meteorological pre-processor), SMOKE (emission pre-processor) and CMAQ (Chemical Transport Model). The meteorological input to the System is the NCEP Global Forecast data. At this stage, the emission input exploits the high resolution inventory for year 2005 produced by TNO, The Netherlands. The Bulgarian national emission inventory for 2010 is used as well. The system is realized on 3 nested domains with increasing resolution.

At the moment, the system is running automatically once a day (00Z). The forecast period is 3 days (72 hours). The results of each System’s run are post-processed in a way to archive the most important pollutants. Part of these pollutants is visualized as sequences of maps giving the evolution of the air quality over Bulgaria and can be seen on the System’s web-site (<http://www.niggg.bas.bg/cw3/>).

ACKNOWLEDGEMENTS

This study is made under the financial support of Bulgarian National Science Fund (Grant Nr. Д002-161/16.12.2008 and ДЦБП-02/1/29.12.2009). The presented results would not be possible without the experience obtained during the participation in the FP5 project BULAIR, the FP6 Network of Excellence ACCENT, the FP6 Integrated Project QUANTIFY, the FP7 projects PASODOBLE (Project No. 241557) and EGI-InSpire (Project No. 261323).

Deep gratitude is due to all organizations providing free of charge data and software used in this study, namely US EPA, US NCEP and European institutions like EMEP, EEA, UBA and many others. Special thanks to the Netherlands Organization for Applied Scientific Research (TNO), for providing the high-resolution European emission inventory, temporal and vertical profiles. Without the open source data bases of these organizations the realization of this system would not be possible.

REFERENCES

- Builtjes, P.J.H., van Loon, M., Schaap, M., Teeuwisse, S., Visschedijk, A.J.H. and Bloos, J.P. (2003) Project on the modelling and verification of ozone reduction strategies: contribution of TNO-MEP, TNO-report, MEP-R2003/166, Apeldoorn, The Netherlands.
- Byun, D. and J. Ching (1999) Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System. EPA Report 600/R-99/030, Washington DC.
- Byun D. and K.L. Schere (2006) Review of the Governing Equations, Computational Algorithms, and Other Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, Applied Mechanics Reviews 59: 51-77.
- CEP (2003): Sparse Matrix Operator Kernel Emission (SMOKE) Modeling System, University of North Carolina, Carolina Environmental Programs - CEP, Research Triangle Park, North Carolina.
- Chen, F., and J. Dudhia, 2001: Coupling an advanced land-surface/ hydrology model with the Penn State/ NCAR MM5 modeling system. Part I: Model description and implementation. *Mon. Wea. Rev.*, 129, 569–585.
- Coats, C.J., Jr., and Houyoux, M.R. (1996): Fast Emissions Modeling With the Sparse Matrix Operator Kernel Emissions Modeling System, The Emissions Inventory: Key to Planning, Permits, Compliance, and Reporting, Air and Waste Management Association, New Orleans, September
- Denier van der Gon, H., A. Visschedijk, H. van de Brugh, R. Dröge (2010) A high resolution European emission data base for the year 2005, TNO-report TNO-034-UT-2010-01895_RPT-ML, Apeldoorn, The Netherlands.
- Dennis, R.L., Byun, D.W., Novak, J.H., Galluppi, K.J., Coats, C.J., and Vouk, M.A (1996): The Next Generation of Integrated Air Quality Modeling: EPA's Models-3, *Atmospheric Environment* 30:1925–1938.
- Dudhia, J., 1989: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model. *J. Atmos. Sci.*, 46, 3077-3107
- EMEP/CORINAIR (2002) Atmospheric emission inventory guidebook, third edition, European Environmental Agency (<http://reports.eea.europa.eu/EMEPCORINAIR3/en/page002.html>).
- European Parliament (2002) DIRECTIVE 2002/3/EC of 12 February 2002 relating to ozone in ambient air, Official Journal of the European Communities (9.3.2002) L67: 14-30.
- Hong, S.-Y., and J.-O. J. Lim, 2006: The WRF Single-Moment 6-Class Microphysics Scheme (WSM6), *J. Korean Meteor. Soc.*, 42, 129–151.
- Hong, S.-Y., and Y. Noh, and J. Dudhia, 2006: A new vertical diffusion package with an explicit treatment of entrainment processes. *Mon. Wea. Rev.*, 134, 2318–2341.
- Houyoux, M.R., and Vukovich, J.M. (1999): Updates to the Sparse Matrix Operator Kernel Emission (SMOKE) Modeling System and Integration with Models-3, The Emission Inventory: Regional Strategies for the Future, Raleigh, NC, Air and Waste Management Association.
- Kain, J. S., 2004: The Kain-Fritsch convective parameterization: An update. *J. Appl. Meteor.*, 43, 170–181.
- Kukkonen J., T. Balk, D. M. Schultz, A. Baklanov, T. Klein, A. I. Miranda, A. Monteiro, M. Hirtl, V. Tarvainen, M. Boy, V.-H. Peuch, A. Poupkou, I. Kioutsioukis, S. Finardi, M. Sofiev, R. Sokhi, K. Lehtinen, K. Karatzas, R. San José, M. Astitha, G. Kallos, M. Schaap, E. Reimer, H. Jakobs, and K. Eben (2012) Operational, regional-scale, chemical weather forecasting models in Europe. *Atmos. Chem. Phys.*, 12, 1–87, 2012, www.atmos-chem-phys.net/12/1/2012/, doi:10.5194/acp-12-1-2012

- Leeuw, Frank de, M. Wim, (2005) Air Quality and Air Quality Indices: a world apart?, ETC/ACC Technical Paper 2005/5, European Topic Centre on Air and Climate Change, European Environmental Agency.
- Mlawer, E. J., S. J. Taubman, P. D. Brown, M. J. Iacono, and S. A. Clough, 1997: Radiative transfer for inhomogeneous atmosphere: RRTM, a validated correlated-k model for the longwave. *J. Geophys. Res.*, 102 (D14), 16663–16682.
- Monteiro, A., Lopes, M., Miranda, A.I., Borrego, C. and Vautard, R. (2005) Air pollution forecast in Portugal: a demand from the new air quality framework directive, *Internat. J. Environment and Pollution* 25: 4–15.
- Poupkou, A., I. Kioutsioukis, I. Lisaridis, K. Markakis, T. Giannaros, E. Katragkou, D. Melas, C. Zerefos and L. Viras (2008a) Evaluation in the Greater Athens Area of an air quality forecast system, *Proc. of the IX EMTE National-International Conference of Meteorology-Climatology and Atmospheric Physics*, 28-31 May 2008, Thessaloniki, Greece, 759-766.
- Poupkou, A., I. Kioutsioukis, I. Lisaridis, K. Markakis, D. Melas, C. Zerefos and Giannaros, T. (2008b): Air quality forecasting for Europe, the Balkans and Athens, 3rd Environmental conference of Macedonia, 14-17 March 2008, Thessaloniki, Greece.
- San Jose, R., J. Perez and R. Gonzalez (2006) Air Quality Real-Time Operational Forecasting System for Europe: an application of MM5-CMAQ-EMIMO modeling system, in Brebia, C. (Ed.), “Air Pollution XIV”, WIT Press, Southampton, Boston, pp. 75-83.
- Schwede, D., G. Pouliot, and T. Pierce (2005): Changes to the Biogenic Emissions Inventory System Version 3 (BEIS3), *Proc. of 4th Annual CMAS Models-3 Users's Conference*, September 26-28, 2005, Chapel Hill, NC.
- Skamarock et al., 2007, “A description of the Advanced Research WRF Version 2”, http://www.mmm.ucar.edu/wrf/users/docs/arw_v2.pdf
- Sofiev, M., Galperin, M., Genikhovich, E. (2008) Construction and evaluation of Eulerian dynamic core for the air quality and emergency modeling system SILAM. NATO Science for piece and security Series C: Environmental Security. Air pollution modelling and its application, XIX, Borrego, C., Miranda, A.I. (eds.), Springer, pp. 699-701.
- Sofiev, M., P. Siljamo, I. Valkama, M. Iivonen, J. Kukkonen (2006) A dispersion modeling system SILAM and its evaluation against ETEX data, *Atmospheric Environment* 40: 674–685.
- Stauffer, D.R. and N.L. Seaman (1990) Use of four-dimensional data assimilation in a limited area mesoscale model. Part I: experiments with synoptic data. *Monthly Weather Review* 118: 1250-1277.
- Syrakov D., K. Ganev, M. Prodanova, K. Slavov, I. Etropolska, N. Miloshev, and G. Jordanov, (2009) Background pollution forecast over Bulgaria, *Journal of International Scientific Publications ECOLOGY & SAFETY*, Volume 3, Part 1 (<http://www.science-journals.eu>), ISSN: 1313-2563, 32-41.
- Syrakov D, I. Etropolska, M. Prodanova, K. Ganev, N. Miloshev, K. Slavov (2012) Operational Pollution Forecast for the Region of Bulgaria, *AIP Conf. Proc.* 1487, 88 (2012); doi: 10.1063/1.4758945.
- Vira J., M. Sofiev (2012) On variational data assimilation for estimating the model initial conditions and emission fluxers for the short-term forecasting of SO_x concentrations. *Atmosph. Environ.*, 46, 318-328, doi:10.1016/j.atmosenv.2011.09.066.